

is related directly to the size (the larger her size, the greater the number of eggs) and age of the female.

#### **Habitats**

Mississippi River; interior rivers and streams; bottomland forests

#### **Iowa Status**

threatened: native

lowa is along the northern extreme of its range. It is associated with backwater areas along large rivers.

# Iowa Range

southeast corner of lowa

### **Bibliography**

Iowa Department of Natural Resources. 2001. Biodiversity of Iowa: Aquatic Habitats CD-ROM.

## diamondback water snake

Nerodia rhombifera

Kingdom: Animalia

Division/Phylum: Chordata - vertebrates

Class: Reptilia
Order: Squamata
Family: Colubridae

### **Features**

The diamondback water snake is 30 to 48 inches long. Its back has a pattern of dark chainlike markings on a background of lighter brown or dirty yellow. The yellow belly is marked with black or brown spots. The scales are keeled (ridged). It is one of the largest of lowa's water snakes.

### **Natural History**

The diamondback water snake lives in lakes, rivers, ditches, ponds, and sloughs. It eats fish and amphibians. It is aquatic, but may be seen basking in the sunshine on limbs over water. It is essentially nocturnal in the hot summer months, but individuals may be active during the day in spring and fall. This snake may flatten its body when disturbed and/or release large amounts of nasty-smelling musk from glands at the base of the tail. Mating occurs in spring. Fourteen to 60 young are born to each female in late summer or early fall. The number of young produced